



| 2024–25 Moderna COVID-19 Vaccine | Ages               | Dose/Injection Amount | Route                        |
|----------------------------------|--------------------|-----------------------|------------------------------|
| Manufacturer-filled syringe      | 5 through 11 years | 0.25 mL/25 $\mu$ g    | Intramuscular (IM) injection |
| Manufacturer-filled syringe      | 12 years and older | 0.50 mL/50 $\mu$ g    | Intramuscular (IM) injection |

### Purpose

To reduce morbidity and mortality from coronavirus disease 2019 (COVID-19) by vaccinating persons who meet the criteria established by the Centers for Disease Control and Prevention's Advisory Committee on Immunization Practices (ACIP).

### Policy

Where authorized under state law, standing orders enable eligible nurses and other health care professionals (e.g., pharmacists) to assess and vaccinate persons who meet the criteria in the "Procedure" section below without the need for clinician examination or direct order from the attending provider at the time of the interaction.

### Procedure

Assess people 5 years of age and older for vaccination with the 2024–25 Moderna COVID-19 Vaccine based on the following criteria:

#### People who are NOT moderately or severely immunocompromised\*

| COVID-19 vaccination history is:   | Administer:  |
|--|--|
| Unvaccinated   | Give 1 dose now.   |
| Any number of previous doses of COVID-19 vaccine, <b>NOT</b> including at least 1 dose of 2024–25 COVID-19 vaccine | Give 1 dose at least 8 weeks after the previous dose. <sup>†</sup> |
| Any number of previous doses COVID-19 vaccine, <b>INCLUDING</b> at least 1 dose of 2024–25 COVID-19 vaccine        | No further doses are indicated.                                    |

\* Persons with a recent SARS-CoV-2 infection may consider delaying vaccination by 3 months from symptom onset or positive test (if infection was asymptomatic).

† For children who turn 5 years of age during the initial series, administer 1 dose of age-appropriate vaccine. If the child has received:

- 1 dose of Moderna COVID-19 vaccine: Administer 1 dose (0.25 mL/25  $\mu$ g) at least 4–8 weeks after Dose 1; there is no dosage change. No further doses are indicated.


**People who ARE moderately or severely immunocompromised**

| COVID-19 vaccination history is:   | Administer:*  |
|--|---|
| Unvaccinated   | Give a 3-dose initial series. Administer:†<br><ul style="list-style-type: none"> <li>■ Dose 1 now</li> <li>■ Dose 2 at least 4 weeks after Dose 1</li> <li>■ Dose 3 at least 4 weeks after Dose 2</li> </ul>  |
| 1 previous dose of any Moderna COVID-19 Vaccine (Dose 1)   | Complete series. Administer:†‡<br><ul style="list-style-type: none"> <li>■ Dose 2 at least 4 weeks after Dose 1</li> <li>■ Dose 3 at least 4 weeks after Dose 2</li> </ul>  |
| 2 doses of any Moderna COVID-19 Vaccine (Doses 1 and 2)  | Complete series. Administer:†‡<br><ul style="list-style-type: none"> <li>■ Dose 3 at least 4 weeks after Dose 2</li> </ul>  |
| 3 or more doses of any mRNA COVID-19 Vaccine, <b>NOT</b> including at least 1 dose of 2024-25 COVID-19 vaccine | Give 1 dose at least 8 weeks after the previous dose.   |
| 3 or more doses of any mRNA COVID-19 Vaccine, <b>INCLUDING</b> at least 1 dose of 2024–25 COVID-19 vaccine     | <ul style="list-style-type: none"> <li>■ People who are moderately or severely immunocompromised may receive 1 additional dose at least 8 weeks following the last recommended dose.</li> <li>■ Further additional dose(s) may be administered, informed by the clinical judgement of a health care provider and personal preference and circumstances.</li> <li>■ Any further additional doses should be administered at least 8 weeks after the last COVID-19 dose.</li> </ul>        |
| 2 or more doses of Novavax, <b>NOT</b> including at least 1 dose of 2024–25 COVID-19 vaccine                   | Give 1 dose at least 8 weeks after the previous dose.   |
| 2 or more doses of Novavax, <b>INCLUDING</b> at least 1 dose of 2024–25 COVID-19 vaccine                       | <ul style="list-style-type: none"> <li>■ People who are moderately or severely immunocompromised may receive 1 additional dose at least 8 weeks following the last recommended dose.</li> <li>■ Further additional dose(s) may be administered, informed by the clinical judgement of a health care provider and personal preference and circumstances.</li> <li>■ Any further additional doses should be administered at least 8 weeks after the last COVID-19 vaccine dose</li> </ul> |

\* People who are moderately or severely immunocompromised should receive the same vaccine product to complete the recommended 3-dose series.

† In the following circumstances, a different age-appropriate COVID-19 vaccine product may be administered: the same vaccine is not available at the time of the clinic visit, the previous dose is unknown, the person would otherwise not receive a recommended dose, or the person starts but is unable to complete a vaccination series with the same vaccine due to a contraindication. For people who received Dose 1 from one manufacturer but will receive subsequent dose(s) from a different manufacturer, administer:

- Dose 2 at least 4 weeks after Dose 1.
- Dose 3 at least 4 weeks after Dose 2.

‡ People should receive the age-appropriate vaccine product and dosage based on their age on the day of vaccination. For children who turn:

- 5 years of age during the 3-dose series should complete the series with the age-appropriate dosage 0.25 mL /25 µg. There is no dosage change.
- 12 years of age during the 3-dose series should complete the series with the age-appropriate dosage 0.5 mL/50µg.

# 2024–2025 Formula Moderna COVID-19 Vaccine

## Standing Orders for Administering Vaccine



### Additional Clinical Considerations

- 2024–25 Moderna COVID-19 Vaccine may be simultaneously administered with other routinely recommended vaccines. There are additional considerations for simultaneous administration of an [orthopoxvirus](#) vaccine and COVID-19 vaccine.
  - People, particularly adolescent or young adult males, who are recommended to receive both vaccines might consider waiting 4 weeks between vaccines.
  - However, if a patient's risk for mpox or severe disease due to COVID-19 is increased, administration of mpox and COVID-19 vaccines should not be delayed.
- Persons who have received hematopoietic cell transplant (HCT) or chimeric antigen receptor (CAR)-T-cell therapy
  - Revaccinate persons who received doses of COVID-19 vaccine prior to or during HCT or CAR-T-cell therapy, following the current COVID-19 vaccination schedule. Revaccination should start at least 3 months (12 weeks) after transplant or CAR-T-cell therapy.
- For additional details and all clinical considerations, see [Interim Clinical Considerations for Use of COVID-19 Vaccines](#).

### Contraindications:

History of a severe allergic reaction (e.g., anaphylaxis) after a previous dose or to a component of the COVID-19 vaccine

### Precautions:

History of:

- A diagnosed non-severe allergy to a component of the COVID-19 vaccine
- Non-severe, immediate (onset less than 4 hours) allergic reaction after administration of a previous dose of one COVID-19 vaccine type, if receiving the same vaccine type
- Moderate to severe acute illness, with or without fever
- Multisystem inflammatory syndrome in children (MIS-C) or adults (MIS-A)
- Myocarditis or pericarditis within 3 weeks after a dose of any COVID-19 vaccine

### Administration

Provide appropriate information material:

- **For 5 through 11 years of age:** Provide all recipients and/or parents/legal guardians with a copy of the current [Fact Sheet for Recipients and Caregivers](#).
- **For 12 years of age and older:** Provide all recipients and/or parents/legal guardians with a copy of the current [COVID-19 Vaccine Information Statement](#).
- Prepare to administer the vaccine following the manufacturer's guidance:
  - [EUA fact sheet for children 5-11 years of age](#)
  - [SPIKEVAX package insert \(fda.gov\)](#)
- Choose the correct [needle gauge, needle length](#), and injection site for persons:
  - **5 through 18 years of age:**
    - » Needle gauge/length: 22–25 gauge, 5/8\*–1-inch
    - » Site: Deltoid muscle of arm<sup>†</sup>
  - **19 years of age and older:** See chart below.
- Administer Moderna COVID-19 Vaccine by intramuscular (IM) injection. Dosage based on age, administer:
  - **5 through 11 years:** 0.25 mL/25 µg
  - **12 years and older:** 0.5 mL/50 µg

\* A 5/8-inch needle can be used if the skin is stretched tightly, and subcutaneous tissues are not bunched.

† Alternately, the anterolateral thigh can be used. A 1- or 1.5-inch needle may be used if administering vaccine in this site, depending on the age of the patient.

# 2024–2025 Formula Moderna COVID-19 Vaccine

Standing Orders for Administering Vaccine



| Sex and Weight of Patient        | Needle Gauge | Needle Length        | Injection Site*       |
|----------------------------------|--------------|----------------------|-----------------------|
| Female or male less than 130 lbs | 22–25        | 5/8 <sup>†</sup> –1" | Deltoid muscle of arm |
| Female or male 130–152 lbs       | 22–25        | 1"                   | Deltoid muscle of arm |
| Female 152–200 lbs               | 22–25        | 1–1.5"               | Deltoid muscle of arm |
| Male 152–260 lbs                 | 22–25        | 1–1.5"               | Deltoid muscle of arm |
| Female 200+ lbs                  | 22–25        | 1.5"                 | Deltoid muscle of arm |
| Male 260+ lbs                    | 22–25        | 1.5"                 | Deltoid muscle of arm |

\* Alternately, the anterolateral thigh can be used. A 1- or 1.5-inch needle may be used if administering vaccine in this site, depending on the age of the patient.

† Some experts recommend a 5/8-inch needle for men and women who weigh less than 130 pounds. If used, skin must be stretched tightly (**do not bunch subcutaneous tissue**).

## Document Vaccination

Document each recipient's vaccine administration information:

- **Medical record:** The vaccine and the date it was administered, manufacturer, lot number, vaccination site and route, name and title of the person administering the vaccine.
- **Vaccination record for recipient:** Date of vaccination, product name/manufacturer, lot number, and name/location of the administering clinic or health care professional.
- **Immunization information system (IIS):** Report the vaccination to the appropriate state/local IIS.

## Be Prepared to Manage Medical Emergencies

Consider observing persons after vaccination to monitor for allergic reactions and syncope:

- **30 minutes** for persons with:
  - A history of a non-severe, immediate (onset within 4 hours) allergic reaction after a previous dose of one COVID-19 vaccine type, if receiving the same vaccine type
  - A history of a diagnosed non-severe allergy to a component of the COVID-19 vaccine, if receiving the same vaccine type
- **15 minutes:** All other persons

Syncope may occur in association with injectable vaccines. Procedures should be in place to avoid falling injuries and manage syncopal reactions.

Have a written protocol to manage medical emergencies following vaccination.

Health care personnel who are trained and qualified to recognize the signs and symptoms of anaphylaxis as well as administer intramuscular epinephrine should be available at the vaccination location at all times.

## Report Adverse Events to the Vaccine Adverse Event Reporting System (VAERS)

### Children 5–11 years of age

**For COVID-19 vaccines given under an Emergency Use Authorization (EUA),** vaccination providers are required to report to [VAERS](#):

- Vaccine administration errors, whether or not associated with an adverse event.
- Serious adverse events regardless of causality. Serious adverse events per FDA are defined as:
  - Death
  - A life-threatening adverse event
  - Inpatient hospitalization or prolongation of existing hospitalization
  - A persistent or significant incapacity or substantial disruption of the ability to conduct normal life functions.
  - A congenital anomaly/birth defect
  - An important medical event that based on appropriate medical judgement may jeopardize the individual and may require medical or surgical intervention to prevent one of the outcomes listed above.
- Cases of Multisystem Inflammatory Syndrome (MIS) in children and adults

# 2024–2025 Formula Moderna COVID-19 Vaccine

## Standing Orders for Administering Vaccine



- Cases of myocarditis
- Cases of pericarditis
- Cases of COVID-19 that result in hospitalization or death.

### Persons 12 years of age and older

For licensed COVID-19 vaccines healthcare providers are strongly encouraged to report to [VAERS](#):

- Any adverse event that occurs after the administration of a vaccine licensed in the United States, whether or not it is clear that a vaccine caused the adverse event.
- Vaccine administration errors, whether or not associated with an adverse event.

Reporting is also encouraged for any other clinically significant adverse event, even if it is uncertain whether the vaccine caused the event. Information on how to submit a report to is available at [Vaccine Adverse Event Reporting System \(VAERS\) \(hhs.gov\)](#) or by calling 1-800-822-7967.

### For More Information, Please See:

- [Interim Considerations: Preparing for the Potential Management of Anaphylaxis after COVID-19 Vaccination](#)
- [CDC's General Best Practice Guidelines for Immunization, "Preventing and Managing Adverse Reactions,"](#)
- [Medical Management of Vaccine Reactions in Children and Teens in a Community Setting](#)
- [Medical Management of Vaccine Reactions in Adults in a Community Setting](#)

**Note:** For more information/guidance, please contact the immunization program at your state or local health department or the appropriate state body (e.g., state board of medical/nursing/pharmacy practice).

### Standing Orders Authorization

This policy and procedure shall remain in effect for all patients of the TotalWellness  
effective September 2024 until rescinded or until June 30, 2025.

Medical director (or other authorized practitioner)

[Signature] / W. Brendan Hayes, M.D. / 09/19/2024  
Signature Date

Adapted with appreciation from the immunize.org standing orders.